

Quantitative Techniques

Pritilata

This reference book can be useful for
BBA, MBA, B.Com, BMS, M.Com, BCA, MCA
and many more courses for Various Universities



**NEERAJ
PUBLICATIONS**
www.neerajbooks.com

Published by:



NEERAJ PUBLICATIONS

(Publishers of Educational Books)

Sales Office : 1507, 1st Floor,

Nai Sarak, Delhi-110 006

E-mail: info@neerajbooks.com

Website: www.neerajbooks.com

© **Reserved with the Publishers only.**

Typesetting by: Competent Computers

Terms & Conditions for Buying E-Book

- The User must Read & Accept the Terms and Conditions (T&C) carefully before clicking on the accept option for Buying the Online Soft Copy of E-books. Under this Particular Facility you may buy only the Online Soft Copy of E-books, no Hard Copy or Printed Copy shall be provided under this facility.
- These E-Books are valid for 365 days online reading only (From the Date of Purchase) and no kind of Downloading, Printing, Copying, etc. are allowed in this facility as these products are just for Online Reading in your Mobile / Tablet / Computers.
- All the online soft copy E-books given in this website shall contain a diffused watermark on nearly every page to protect the material from being pirated / copy / misused, etc.
- This is a Chargeable Facility / Provision to Buy the Online Soft Copy of E-books available online through our Website Which a Subscriber / Buyer may Read Online on his or her Mobile / Tablet / Computer. The E-books content and their answer given in these Soft Copy provides you just the approximate pattern of the actual Answer. However, the actual Content / Study Material / Assignments / Question Papers might somewhat vary in its contents, distribution of marks and their level of difficulty.
- These E-Books are prepared by the author for the help, guidance and reference of the student to get an idea of how he/she can study easily in a short time duration. Content matter & Sample answers given in this E-Book may be Seen as the Guide/Reference Material only. Neither the publisher nor the author or seller will be responsible for any damage or loss due to any mistake, error or discrepancy as we do not claim the Accuracy of these solution / Answers. Any Omission or Error is highly regretted though every care has been taken while preparing these E-Books. Any mistake, error or discrepancy noted may be brought to the publishers notice which shall be taken care of in the next edition. Please consult your Teacher/Tutor or refer to the prescribed & recommended study material of the university / board / institute / Govt. of India Publication or notification if you have any doubts or confusions before you appear in the exam or Prepare your Assignments before submitting to the University/Board/Institute.
- Publisher / Study Badshah / shall remain the custodian of the Contents right / Copy Right of the Content of these reference E-books given / being offered at the website www.studybadshah.com.
- The User agrees Not to reproduce, duplicate, copy, sell, resell or exploit for any commercial purposes, any portion of these Services / Facilities, use of the Service / Facility, or access to the Service / Facility.
- The Price of these E-books may be Revised / Changed without any Prior Notice.
- The time duration of providing this online reading facility of 365 days may be alter or change by studybadshah.com without any Prior Notice.
- The Right to accept the order or reject the order of any E-books made by any customer is reserved with www.studybadshah.com only.
- All material prewritten or custom written is intended for the sole purpose of research and exemplary purposes only. We encourage you to use our material as a research and study aid only. Plagiarism is a crime, and we condone such behaviour. Please use our material responsibly.
- In any Dispute What so ever Maximum Anyone can Claim is the Cost of a particular E-book which he had paid to Study Badshah company / website.
- If In case any Reader/Student has paid for any E-Book and is unable to Access the same at our Website for Online Reading Due to any Technical Error/ Web Admin Issue / Server Blockage at our Website www.studybadshah.com then He will be send a New Link for that Particular E-Book to Access the same and if Still the Issue is Not Resolved Because of Technical Error/ Web Admin Issue / Server Blockage at our website then His Amount for that Particular Purchase will be refunded by our website via PayTM.
- All the Terms, Matters & Disputes are Subjected to "Delhi" Jurisdiction Only.

CONTENTS

S.No.	Page
1. Functions, Limit and Continuity	1
2. Derivatives	7
3. Partial Differentiation	14
4. Maxima and Minima	20
5. Unconstrained Optimization	26
6. Constrained Optimization	28
7. Integration and Applications in Economic Dynamics	31
8. Difference Equations and Applications in Economic Dynamics	42
9. Vector Analysis	49
10. Linear Algebra	56
11. Input-output Analysis	69
12. Linear Programming	77
13. Data Presentation and Descriptive Statistics	87
14. Correlation and Regression Analysis	101
15. Probability Theory	110
16. Probability Distribution	121
17. Sampling Theory	128
18. Sampling Distribution	134
19. Statistical Inference	142
20. Set Theory	156
21. Functions and their Graphical Representation	173
22. Integration Techniques	190

Sample Preview of The Chapter

Published by:



**NEERAJ
PUBLICATIONS**

www.neerajbooks.com

QUANTITATIVE TECHNIQUES

INTRODUCTION TO DIFFERENTIAL CALCULUS



Functions, Limit and Continuity

INTRODUCTION

Mathematical Techniques are very useful in order to solve economic problems. Some concepts are related to function, limit, and continuity are expressed here. There are many ways to describe or represent functions: by a formula, by an algorithm that computes it, or by plotting graph. Making relationship between describing the problem and then interpreting it, is the main function of these mathematical expressions. In order to clear the concepts we will discuss some illustrations and also some practical exercises. The explanations given here will manage the understanding of different mathematical and statistical knowledge levels of students in each and every aspect.

CHAPTER AT A GLANCE

REVIEW OF THE BASIC CONCEPTS

Set

Sets are “collections”. The objects “in” the collection are its members, e.g. we are all members of the set of all humans. There are sets of numbers, people, and other sets.

Example: $\{x : x \text{ is an even number}\}$

The set containing the even numbers (i.e. $\{0, 2, 4, \dots\}$)

FUNCTION

It is defined as the one quantity (the argument of the function, also known as the input) that completely determines another quantity (the value, or the output).

Domain: The set of all permitted inputs to a given function is called the domain of the function. Range: The set of all resulting outputs is called the image or range of the function.

Co-domain: The image is often a subset of some larger set, called the co-domain of a function. For example, the function $f(x) = x^2$ could take as its domain the set of all real numbers, as its image the set of all non-negative real numbers, and as its co-domain the set of all real numbers.

A function assigns exactly one value to each input of a specified type. The argument and the value may be real numbers, but they can also be elements from any given sets: the domain and the co-domain of the function.

Example of a function with the real numbers as both its domain and co-domain:

Function $f(x) = 2x$, which assigns to every real number the real number with twice its value.

In this case, it is written that $f(5) = 10$.

The notation $f : X \rightarrow Y$ indicates that f is a function with domain X and co-domain Y .

Variable: These are numerical values which changes with different mathematical operations. For example, a, b, x, y, \dots

Continuous Variable: If y takes all prime numbers from a given number a to another given number b , then y is known as a continuous variable.

Interval: The value between two terminal values are known as interval. In the expression $(x + 3)(x + 5)$, the interval of x is -3 and -5 .

Constant

The numerical value which never changes with change in mathematical operation. i.e. $5, 6, \frac{1}{2}\pi, \dots$

Absolute Value: The absolute value of a number is its distance from 0 on a number line.

For example, the number 9 is 9 units away from 0. Therefore its absolute value is 9.

2 / NEERAJ : QUANTITATIVE TECHNIQUES

In case of negative number, the absolute value is also positive. The number -4 is still 4 units away from 0. The absolute value of -4 is therefore positive 4.


Examples:

$$\begin{aligned} |4| &= 4 \\ |-4| &= 4 \\ |4 + 3| &= 7 \\ |-4 - 3| &= 7 \\ |3 - 4| &= 1 \\ -|4| &= -4 \\ -|-4| &= -4 \end{aligned}$$


Graph of a Function

The symbol for the input to a function is often called the independent variable or argument and is often represented by the letter x or, if the input is a particular time, by the letter t . The symbol for the output is called the dependent variable or value and is often represented by the letter y . The function itself is most often called f , and thus the notation $y = f(x)$ indicates that a function named f has an input named x and an output named y .

Graph of the function $(x^2 - 4)/(x - 2)$



Graph of the function $\sin(x)/x$



Bounded Functions and their Bounds

A function f defined on some set X with real or complex values is called bounded, if the set of its values is bounded. In other words, there exists a real number $M < \infty$ such that

$$|f(x)| \leq M \text{ for all } x \text{ in } X.$$

Sometimes, if $f(x) \leq A$ for all x in X , then the function is said to be bounded above by A . On the other hand, if $f(x) \geq B$ for all x in X , then the function is said to be bounded below by B .

Monotone Function

A monotonic function (or monotone function) is a function that preserves the given order. A function f defined on a subset of the real numbers with real values is called monotonic (also monotonically increasing, increasing or non-decreasing), if for all x and y such that $x \leq y$ one has $f(x) \leq f(y)$, so f preserves the order. Likewise, a function is called monotonically decreasing (also decreasing or non-increasing) if, whenever $x \leq y$, then $f(x) \geq f(y)$, so it reverses the order.

If the order \leq in the definition of monotonicity is replaced by the strict order $<$, then one obtains a stronger requirement. A function with this property is called strictly increasing. Again, by inverting the order symbol, one finds a corresponding concept called strictly decreasing. Functions that are strictly increasing or decreasing are one-to-one (because for x not equal to y , either $x < y$ or $x > y$ and so, by monotonicity, either $f(x) < f(y)$ or $f(x) > f(y)$, thus $f(x)$ is not equal to $f(y)$).

Inverse Function

If f is a function from a set A to a set B , then an inverse function for f is a function from B to A , with the property that a round trip (a composition) from A to B to A (or from B to A to B) returns each element of the initial set to itself. Thus, if an input x into the function f produces an output y , then inputting y into the inverse function produces the output x , and vice versa.

Types of Function

1. Algebraic Function: An algebraic function is informally a function that satisfies a polynomial equation whose coefficients are themselves polynomials. For example, an algebraic function in one variable x is a solution y for an equation where the coefficients $a_i(x)$ are polynomial functions of x .

A constant function is a function whose values do not vary and thus are constant. For example,

if we have the function $f(x) = 4$, then f is constant since f maps any value to 4. More formally, a function $f: A \rightarrow B$ is a constant function if $f(x) = f(y)$ for all x and y in A .

2. **Non-Algebraic Function:** A function which is not algebraic is called a transcendental function. i.e. $y = pqx$

CONCEPT OF LIMIT

Limit of a Function

The Limit of $f(x)$ as x approaches a is L :

$$\lim_{x \rightarrow a} f(x) = L$$

Example:

$$\lim_{x \rightarrow -2} f(x) \text{ where } f(x) = \begin{cases} -3x & \text{if } x \neq -2 \\ 1 & \text{if } x = -2 \end{cases}$$

$$\begin{aligned} \lim_{x \rightarrow -2} f(x) &= \lim_{x \rightarrow -2} -3x \\ &= -3 \lim_{x \rightarrow -2} x \\ &= -3(-2) = 6 \end{aligned}$$

Right Hand Limit of a Function

The right-hand limit of $f(x)$, as x approaches a , equals L

$$\lim_{x \rightarrow a^+} f(x) = L$$

Left Hand Limit of a Function

The left-hand limit of $f(x)$, as x approaches a , equals M

$$\lim_{x \rightarrow a^-} f(x) = M$$

Examples of One-Sided Limit

$$\lim_{x \rightarrow 3^+} f(x) = \lim_{x \rightarrow 3^+} 2x = 6$$

$$\lim_{x \rightarrow 3^-} f(x)$$

$$\lim_{x \rightarrow 3^-} f(x) = \lim_{x \rightarrow 3^-} x^2 = 9$$

Functions Tending to Infinity:

If the extended real line R is considered, i.e. $R \cup \{-\infty, \infty\}$, then it is possible to define limits of a function at infinity.

If $f(x)$ is a real function, then the limit of f as x approaches infinity is L , denoted

$$\lim_{x \rightarrow \infty} f(x) = L,$$

if and only if for all $\epsilon > 0$ there exists $S > 0$ such that $|f(x) - L| < \epsilon$ whenever $x > S$.

Similarly, the limit of f as x approaches negative infinity is L , denoted

$$\lim_{x \rightarrow -\infty} f(x) = L,$$

if and only if for all $\epsilon > 0$ there exists $S < 0$ such that $|f(x) - L| < \epsilon$ whenever $x < S$.

For example

$$\lim_{x \rightarrow -\infty} e^x = 0$$

Fundamental Theorems on Limit

If c is any number, $\lim_{x \rightarrow a} f(x) = L$ and $\lim_{x \rightarrow a} g(x) = M$, then

(a) $\lim_{x \rightarrow a} (f(x) + g(x)) = L + M$

(b) $\lim_{x \rightarrow a} (f(x) - g(x)) = L - M$

(c) $\lim_{x \rightarrow a} (f(x) \cdot g(x)) = L \cdot M$

(d) $\lim_{x \rightarrow a} (f(x) / g(x)) = L/M, M \neq 0$

(e) $\lim_{x \rightarrow a} (c \cdot f(x)) = c \cdot L$

(f) $\lim_{x \rightarrow a} (f(x))^n = L^n$

(g) $\lim_{x \rightarrow a} c = c$

(h) $\lim_{x \rightarrow a} x = a$

(i) $\lim_{x \rightarrow a} x^n = a^n$

(j) $\lim_{x \rightarrow a} \sqrt[n]{f(x)} = \sqrt[n]{L}, L > 0$

CONTINUITY

A function f is continuous at the point $x = a$ if the following are true:

(i) $f(a)$ is defined

(ii) $\lim_{x \rightarrow a} f(x)$ exists

If f and g are continuous at $x = a$, then $f \pm g, fg$ and f/g ($g(a) \neq 0$) are continuous at $x = a$

A polynomial function $y = P(x)$ is continuous at every point x .

A rational function $R(x) = P(x)/Q(x)$ is continuous at every point x in its domain.

Given $f(x) = 3x^2 - 2x - 5$,

Show that $f(x) = 0$ has a solution on $[1, 2]$.

4 / NEERAJ : QUANTITATIVE TECHNIQUES

$$f(1) = -4 < 0$$

$$f(2) = 3 > 0$$

$f(x)$ is continuous (polynomial) and since $f(1) < 0$ and $f(2) > 0$, by the Intermediate Value Theorem there exists $a \in c$ on $[1, 2]$ such that $f(c) = 0$.

Limits at Infinity

For all $n > 0$,

$$\lim_{x \rightarrow \infty} \frac{1}{x^n} = \lim_{x \rightarrow -\infty} \frac{1}{x^n} = 0$$

provided that $1/x^n$ is defined.

CHECK YOUR PROGRESS

Q.1. Distinguish between variable and constants giving examples.

Ans. Variable changes with the change in numerical values whereas constant does not change with their numerical changes.

For Example: variables, a, b, c, d, e, \dots

Constant, $\pi, 1/3, 4/9, \dots$

Q. 2. Point out the domain of definition of the following functions:

(i) $(\cos x + \sin x) / (\cos x - \sin x) = f(x)$

Sol. Let $(\cos x + \sin x) / (\cos x - \sin x) = 0$

Domain = Any real value of x except $x = \frac{\pi}{4}$.

(ii) $\sqrt{x^2 - 5x + 6x} = f(x)$

Sol. Let $\sqrt{x^2 - 5x + 6x} = 0$

or, $x^2 - 3x - 2x + 6 = 0;$

or, $x(x - 3) - 2(x - 3) = 0;$

or, $(x - 2)(x - 3) = 0;$

or, $x = 2$ or, $x = 3$

Domain = All real value of x except $2 < x < 3$.

(iii) $f(x) = \sin^{-1} x$

Sol. Let $\sin^{-1} x = 0$,

Value of \sin lies between -1 and 1

Domain = $-1 < x < 1$

(iv) $\log(3x - 1) = f(x)$

Sol. Let $\log(3x - 1) = 0$

or, $3x - 1 = 0$


or, $x = 1/3$

Domain = value of $x > 1/3$.

Q. 3. $q = f(p) = 1/p$

Draw the graph of the above function. Show that this is a monotonically decreasing function. Also, show that the function has a lower bound (zero).

Ans.



Q. 4. $f(x) = x^3 + 2x$. Find $\frac{f(x+h) - f(x)}{h}$.

Sol. $\frac{f(x+h) - f(x)}{h} = \frac{f(x+h)}{h} - \frac{f(x)}{h}$

Putting the value of $(x + h)$ at place of x ,

$$\begin{aligned} \frac{f(x+h)}{h} &= \frac{\{(x+h)^3\}}{h} + \frac{\{2(x+h)\}}{h} \\ &= \frac{(x^3 + h^3 + 3x^2h + 3xh^2)}{h} + \frac{(2x + 2h)}{h} \\ &= \frac{x^3}{h} + \frac{h^3}{h} + 3x^2 + 3xh + \frac{2x}{h} + 2 \\ f(x+h) - f(x) &= \frac{x^3}{h} + \frac{h^3}{h} + 3x^2 + 3xh + \frac{2x}{h} - \frac{2x}{h} \\ &= 3x^2 + 3xh + h^2 + 2. \end{aligned}$$

Q. 5. Evaluate the following limits:

(i) $\lim_{x \rightarrow 0} (\sqrt{a+x} - \sqrt{a-x}) / x$

Sol. $\lim_{x \rightarrow 0} (\sqrt{a+x} - \sqrt{a-x}) / x$

$$= \lim_{x \rightarrow 0} \left(\frac{\sqrt{a+x} - \sqrt{a-x}}{x} \cdot \frac{\sqrt{a+x} + \sqrt{a-x}}{\sqrt{a+x} + \sqrt{a-x}} \right)$$

$$= \lim_{x \rightarrow 0} \left(\frac{(\sqrt{a+x})^2 - (\sqrt{a-x})^2}{x} \cdot \frac{1}{\sqrt{a+x} + \sqrt{a-x}} \right)$$

$$= \lim_{x \rightarrow 0} \left(\frac{(\sqrt{a+x}) - (\sqrt{a-x})}{x} \cdot \frac{1}{\sqrt{a+x} + \sqrt{a-x}} \right)$$

$$= \lim_{x \rightarrow 0} \frac{2x}{x\sqrt{a+x} + \sqrt{a-x}}$$

$$= \lim_{x \rightarrow 0} \left(\frac{2}{\sqrt{a+x} + \sqrt{a-x}} \right)$$